

Sulphaquinoxaline W.S.P

For veterinary use only

Composition:

Each 100 gm contains:

Sulphaquinoxaline Sodium 20 g

(Equivalent to Sulphaquinoxaline base 18.63 g

Properties:

Water soluble Sulphaquinoxaline for veterinary use is a broad Spectrum Chemotherapeutic agent which is very effective against all types of coccidia, beside its antimicrobial activity against pasterella species and some other bacterial infections affecting poultry. Water soluble Sulphaquinoxaline is highly recommended to treat outbreaks of all forms of coccidiosis in chicken, turkeys and rabbits only.

Indications:

- Treatment of coccidiosis in poultry, turkeys and rabbits only.
- Treatment of cholera, typhoid and paratyphoid in poultry, turkeys and rabbits only.

Precautions:

- Sulphonamides may cause petechial haemorrhages in poultry as a result of vitamin K antagonism.
- It may cause toxic reactions unless the drug is evenly mixed in water at doses indicated and used according to direction.
- It is hazardous as prolonged administration at higher doses may result in depressed feed intake, deposition of sulphaquinoxaline crystals in the kidney and interference with normal blood clotting.
- Also prolonged treatment may lead to vitamin K deficiency, agranulocytosis and haemolytic anemia.
- It causes skin and eye burns so avoid contact with eye, skin or clothing. In case of contact, flush immediately with water for at least 15 minutes.
- It is used as directed due to its restrictions.

Dosage and administration:

Poultry and Turkeys:

- Treatment of coccidiosis:
3 g Powder of product per liter of drinking water (equivalent to 500 mg of active ingredient per liter of drinking water) for 3 consecutive days

(the solution should be stirred well), rest for 3 days with plain drinking water then repeat the treatment with a dose of 1 gm / liter for 2 more days.

- Treatment of fowl cholera, typhoid and paratyphoid:

3 g. per liter of drinking water (equivalent to 500 mg of active ingredient per liter of drinking water) for 5 consecutive days.

Rabbits:

- Treatment of coccidiosis:

3 g per liter of drinking water (equivalent to 500 mg of active ingredient per liter of drinking water) for 2 weeks.

Withdrawal time:

- Withdrawal time: 10 days

- It should not be used in chicken and turkeys producing eggs for human consumption.

Storage:

Store in a cool dry place protected from light and heat exposure.

Note:

Opened packs should be used promptly

Packing:

500 g and 1 Kg aluminium foil bags.